

ORIGINAL ARTICLE

PREVALENCE OF ANXIETY AND DEPRESSION AMONG MEDICAL STUDENTS IN MMCH, SINDH, PAKISTAN

*Razzaq S A, **Asghar H, ***Irshad H A, ****Samoon N A, *****Muhammad S R

ABSTRACT

Objectives:

To know the prevalence of anxiety and depression among medical students of Muhammad Medical College, Mirpurkhas, Sindh, Pakistan

Methods:

A Cross-sectional study was done on the students of Muhammad Medical College, Mirpurkhas, Sindh, who had spent more than six months in the medical college. Self administered questionnaire were given to the students, present in the class and willing to participate in the study

The instrument used to assess the anxiety and depression levels was the, Aga Khan University Anxiety and Depression Scale (AKUADS). Additional questions regarding socioeconomic variables were also included in the survey instrument, such as student's age order, family monthly income, and gender, marital status.

Results:

Out of 380 students, 330 completed the questionnaire from final year MBBS to 1st year MBBS. The 330 students present in survey among them 108 (32.72%) were males and 222 (67.27%) females students.

Using anxiety and depression scale it was found out that 318 (96.36%) students had anxiety and depression. Among them 215 (67.61%) were females and 103 (32.38%) males.

Prevalence of anxiety and depression in students of Final, 4th year and 3rd year 2nd and 1st years was 13%, 15%, 20% and 21%, 28% respectively. It was significantly higher in 1st year and 2nd year students, as compared to final and 4th year and students.

Using AKUADS, it was found that 47.93% of students suffered from anxiety and depression. Anxiety and depression was slightly higher in 1st year MBBS students and 2nd year MBBS students & lower in Final year and 4th year MBBS students. It was also higher in female students as compared to male students.

CONCLUSIONS:

The result shows that most of medical students have anxiety and depression due to study burden and a busy schedule.

It is also seen that anxiety and depression is more among female medical students, and more among in 1st year and 2nd year MBBS students, as compared to the final and forth year students.

Keywords: Anxiety, depression, Medical students, prevalence

Introduction:

Medical education is stressful. High levels of stress have been seen in medical students in various studies. (1-4) among medical students, anxiety and depression has been reported to be due to their academic stressors include the volume of material to be learned, academic performance and evaluation (examination and continuous assessment, mental tension and too much work load. (5) All this can result in decreased life satisfaction among students. (6) Stress during medical school can lead to problems later in professional life compromising patient care. (7) Anxiety is defined as excessive worry and tension, on most days, for at least 6 months, together with the following sign and symptoms; increase motor tension (fatigue, trembling, restlessness, and muscle tension); autonomic hyperactivity (shortness of breath, rapid heart rate, dryness of mouth, cold hand and dizziness). If not detected early and treated anxiety lead to lifelong problems and has co-morbidity with depression. Anxiety causes disruption on day to day functioning and reduction in quality of life. (8) A study among British medical students found that 39% had clinically significant levels of anxiety. (9) Depression is one of the most prevalent mental disorders today, and it is characterized by episodes of low mood associated with loss of interest in daily activities. According to the World Health Organization (WHO), depression is the third most common disabling disorder, especially among females. (10) Depression and anxiety have been reported universally among medical students as they are exposed to stressors. (11, 12, 13) This situation can lead to mental distress and have adverse effects on their cognitive functioning. (14, 15) Many authors mention that medical students have

Elevated levels of depression and anxiety compared to those who are not in the medical field. (16) However, numerous studies have been conducted about the factors related to depression and anxiety in medical students. Some of these factors include academic load, insomnia, watching patient's suffering, and financial concerns. (17) This study may help in designing appropriate intervention strategies to enhance the learning abilities which will improve the patient care by achieving good communication, increasing quality of care and decreasing medical errors in addition to burden the costs paid by the society through anxiety and depression attenuation.

Materials and methods:

A cross sectional study was done on students of Muhammad Medical College, Mirpurkhas, Data collection spanned over the month of May 2013. The approval was obtained from the principle and Dean of the college before administering questionnaires. Verbal consent was taken from students before distributing questionnaires. Performance and confidentiality was ensured. The students who had spent more than six months in the medical college were included in this study, the survey was done at a time when the students did not have any major examination scheduled.

The instrument used to assess the anxiety and depression levels was, Aga Khan University Anxiety and Depression Scale (AKUADS). (18) It has been validated on a statistically appropriate sample size, in urban squatter settlement of Karachi. (19) At a cut off score of 19 points KUADS has specificity of 81%, sensitivity of 74%, a positive predictive value of 63%, and negative predictive value of 88%. (20)

which is higher than other available scales like the self-reporting questionnaire (SRQ). (21, 22) The students were instructed to return the completed questionnaire. Out of 517 students, 380 were present during the survey. We distributed the forms which were returned later on. A total of 330 out of 380 distributed forms were received with a response rate of 86.84%.

To determine the prevalence of anxiety and depression, more questions were also included in the survey instrument, such as student's age, gender, locality, marital status, family monthly income, and monthly expenditure on education. To ensure anonymity, the respondents were asked not to put names or other identifying notation on the questionnaire.

Results:

There were total 517 students from final year MBBS to 1st year MBBS.

In our survey there is 380 students present but questionnaire respond were given by 330. The 330 students present in survey among them 108 (32.72%) were males and 222 (67.27%) females students. Using anxiety and depression scale it was found out that 318 (96.36%) students had anxiety and depression. Among them 215 (67.61%) were females and 103 (32.38%) males.

Prevalence of anxiety and depression in students of final, 4th year and 3rd year 2nd and 1st years was 13%, 15%, 20% and 21%, 28% respectively.

It was significantly higher in 1st year and 2nd year students, as compared to final and 4th year and students.

Table-2 gives the frequency of students in different groups with respect to

Table 1: Anxiety and Depression causes according to class and gender

Batch	Number of students			Anxiety and depression		
	Male	Female	Total	Male No. & %	Female No. & %	Total No. %
Final Year	18	36	54	19(18.44)	24(11.16)	43(13.52)
4 th year	12	49	61	18(17.47)	32(14.88)	50(15.72)
3 rd year	25	39	64	22(21.35)	44(20.46)	66(20.75)
2 nd year	21	52	73	19(18.44)	50(23.25)	69(21.69)
1 st year	32	46	78	25(24.27)	65(30.23)	90(28.30)
Total	108	222	330	103(32.38)	215(67.61)	318(96.36)

Table 2:

Association of anxiety & depression with demographic and social factors

Variable	n.	% 100%	Anxiety and depression	
			present	absent
Total participants	330	100%		
Age				
< 20 Years	105	31.81%	55	50
22-24 Years	150	45.45%	62	88
> 24 Years	75	22.72%	24	51
Gender				
Male	108	32.72%	36	72
Female	222	67.27%	126	96
Marital Status				
Married	28	8.48%	16	12
Unmarried	302	91.51%	145	157
Locality				
Urban	284	86.0%	122	162
Rural	46	13.93%	19	27
Total Family Income				
< Rs. 25,000	74	22.42%	36	38
Rs.15,000-Rs.30,000	94	28.48%	54	40
Rs.30,000-Rs.50,000	84	25.45%	39	45
> Rs.50,000	80	24.24%	35	45

Table-2 gives the frequency of students in different groups with respect to age, gender, marital status, locality, total family income. Using AKUADS, it was found that 47.93% of students suffered from anxiety and depression. Anxiety and depression was slightly higher in 1st year MBBS students and 2nd year MBBS students & lower in Final year and 4th year MBBS students. It was also higher in female students as compared to male students. It was seen that higher in unmarried.

Discussion:

WHO estimates that one in five people are suffering from depression and anxiety. (23) Many studies reported significant distress in medical students (24, 25, 26) Based on the medical students in our study, the prevalence of anxiety and depression was 47.9% in MMCH, Sindh. Also in our study shows the percentage ration of female medical students are higher than male students due to self expectation, feeling of lack of competence and tendency to over report symptoms. that 67.27% in female and 32.72% in male students.

The results of our study different significantly from some other studies due to

the difference in teaching and assessment methodologies including introduction of problem based learning and objective structured performance evaluation in the recent years. In our study one thing in Difference from other studies is that percentage of anxiety and depression is 47.9% the difference is sample size of students were 330 out of 517 total students.

In our study anxiety and depression decreased with increasing age. This may be due to adapt to stress of clinical training after spending a years in it. Anxiety and depression can lead to negative outcomes including medical school dropout, not to do work properly, not having good relationships with friends, burnout, and increased suicidal tendency.

There is need for greater attention for medical students to psychological point of view.

It is necessary for medical student to seek appropriate help for mental health problem and view it as a weakness. Medical college should encourage the stressful students to spend adequate time on their social and personal lives and emphasize the importance of health promoting coping strategies.

Preventive programming should begin in medical colleges to address the academic and financial and personal problems, early sign and symptoms of anxiety and depression should be addressed.

We need strategy for students to cope with stress to make a smooth transition life from medical college towards professional life and also to adjust to different learning environments during different phases of medical education.

It is also necessary that base line information should be given to the medical students at the time of entrance in medical colleges. And further evaluation of positive cases should be done by a Psychiatrist to improve mental health of students. Follow up studies for monitoring prevalence of anxiety and depression will help in instituting interventional strategies.

It is noted that proportion of medical students had ongoing psychiatric conditions. Action should be taken to find out psychological problems and to provide adequate facilities to the stressful students. Medical colleges also courage the stressful student from the beginning (from 1st year) to relieve depressive symptoms

Conclusion:

From our study, it is concluded that most of medical students have anxiety and depression then from other professional due to hard work load.

It is also seen that anxiety and depression is more among female medical students, and more among in 1st year and 2nd year MBBS students, as compared to the final and forth year students

Acknowledgment:

We Thank Prof. Dr. Syed Razi Muhammad and Dr. Noor Ali Samoon for his encouragement and permission to publish

References:

1. The prevalence and socio-demographic correlations of depression, anxiety and stress among a group of university students. Soc Psychiatry Psychiatr Epidemiol 2008; 43: 667-72.
2. Gale C, Oakley-Bronme M. Anxiety disorder. BMJ 2003;321 : 1204-1207;
3. Karaman T,Y, Erdem S, Erengin H, Akaydin M. Anxiety, depression and stressful life events among medical students: a prospective study in Antalya, Turkey. Med Educ 2001; 35: 12-7.

4. Azhar MZ. Psychological stress and treatment - research issues. *Med J Malaysia* 2004; 59: 143-5.
5. Cozens J. Medical student stress. *Med Educ* 2001, 2006; 35: 6-7.
6. Arnold L, et al. Medical students perception of stress, gender and ethnic consideration. *Acad Med* 1994;69(Supp. I):S22-4.
7. Gronvold NT, et al. Life satisfaction and resilience in medical school--a six-year longitudinal, nationwide and comparative study. *BMC Med Educ* 2006; 6: 48.
8. Ekeberg O. Factors in medical school that predict postgraduate mental health problems in need of treatment. A nationwide and longitudinal study. *Med Educ* 2001; 35: 110-20.
9. Ashton CH, Kamali F. Personality, lifestyles, alcohol and drug consumption in a sample of British medical students. *Medical education* 1995;29; 187-192
10. Ibrahim N, Al-Kharboush D, El-Khatib L, Al-Habib A, Asali D. Prevalence and Predictors of Anxiety and Depression among Female Medical Students in King Abdulaziz University, Jeddah, Saudi Arabia. *Iran J Public Health* [Internet]. 2013 [cited 2016 Dec 18];42(7):726–36.
11. Ahmed I, Banu H, Al-Fageer R, Al-Suwaidi R. Cognitive emotions: Depression and anxiety in medical students and staff. *J Crit Care* [Internet]. 2009 Sep [cited 2016 Dec 18];24(3):e1–7.
12. Sohail N. Stress and academic performance among medical students. *J Coll Physicians Surg Pak* [Internet]. 2013 Jan [cited 2016 Dec 18];23(1):67–71.
13. Abdulghani HM, AlKanhah AA, Mahmoud ES, Ponnampereuma GG, Alfaris EA. Stress and its effects on medical students: a cross-sectional study at a college of medicine in Saudi Arabia. *J Health Popul Nutr* [Internet]. 2011 Oct [cited 2016 Dec 18];29(5):516–22.
14. Dahlin M, Joneborg N, Runeson B. Stress and depression among medical students: a cross-sectional study. *Med Educ* [Internet]. 2005 Jun [cited 2016 Dec 18];39(6):594–604.
15. Baldassin S, Alves TC de TF, de Andrade AG, Nogueira Martins LA. The characteristics of depressive symptoms in medical students during medical education and training: a cross-sectional study. *BMC Med Educ* [Internet]. 2008 Dec 11 [cited 2016 Dec 18];8(1):60.
16. Dyrbye LN, Thomas MR, Shanafelt TD. Systematic review of depression, anxiety, and other indicators of psychological distress among U.S. and Canadian medical students. *Acad Med* [Internet]. 2006 Apr [cited 2016 Dec 18];81(4):354–73.
17. Roh M-S, Jeon HJ, Kim H, Han SK, Hahm B-J. The Prevalence and Impact of Depression Among Medical Students: A Nationwide Cross-Sectional Study in South Korea. *Acad Med* [Internet]. 2010 Aug [cited 2016 Dec 18];85(8):1384–90.
18. All B, Hashim RM, Khan MM, et al. Development of an indigenous screening instrument in Pakistan: the Aga Khan University anxiety and depression scale. *J Pak Med Assoc* 1998;48:261-5.
19. All B. Validation of an indigenous screening questionnaire for anxiety and depression in an urban squatter settlement of Karachi, *J Coll Physician Surg* 1998; 8:207-10.
20. Ali B, Raza H, Khan MM, Jehan I. Development of an indigenous screening instrument in Pakistan: the Aga Khan University Anxiety and Depression scale. *J Pak Med Assoc* 1998; 48: 261-5.
21. Ali BS, Amanullah S. A comparative review of two screening instruments: the Aga Khan University anxiety and depression scale and the self reporting questionnaire. *J Pak Med Assoc* 1998; 48: 79-82.
22. Ahmer S, Faruqui RA, Aijaz A. Psychiatric rating scales in Urdu: a systematic review. *BMC Psychiatry* 2007; 7: 59.
23. WHO | Investing in treatment for depression and anxiety leads to fourfold return. WHO. 2016;
24. Clark DC, Zeldow PB. Vicissitudes of depressed mood during four years of medical school. *JAMA* [Internet]. 1988 Nov 4 [cited 2016 Dec 18];260(17):2521–8.
25. Vitaliano PP, Maiuro RD, Russo J, Mitchell ES, Carr JE, Van Citters RL. A biopsychosocial model of medical student distress. *J Behav Med* [Internet]. 1988 Aug [cited 2016 Dec 18];11(4):311–31.
26. Inam SNB, Saqib A, Alam E. Prevalence of anxiety and depression among medical students of private university. *J Pak Med Assoc* [Internet]. 2003 Feb [cited 2016 Dec 18];53(2):44–7.

*Syed Abdul Razzaq

Student of Muhammad Medical College, Mirpurkhas

**Hassan Asghar

Student of Muhammad Medical College, Mirpurkhas

***Hafiz Aqeel Irshad

Student of Muhammad Medical College, Mirpurkhas

****Dr. Noor Ali Samoon

Assistant Professor of Community Medicine, Muhammad Medical College, Mirpurkhas

*****Prof. Dr. Syed Razi Muhammad

Dean and Professor of Surgery Muhammad Medical College, Mirpurkhas